Anti-CCR2 antibody(DMC447); IgG1 Chimeric mAb

Cat. No. DMC100447

## PRODUCT INFORMATION

| | |
|---|---|
| **Clone ID** | **Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** DMC447 |
| **Target** | CCR2 |
| **Synonyms** | CC-CKR-2; MCP-1-R; CD192 |
| **Host Species** | Rabbit |
| **Description** | Anti-CCR2 antibody(DMC447); IgG1 Chimeric mAb |
| **Delivery** | In Stock |
| **Uniprot ID** | P41597 |
| **IgG type** | Rabbit/Human Fc chimeric IgG1 |
| **Clonality** | Monoclonal |
| **Reactivity** | Human |
| **Applications** | Flow Cyt |
| **Recommended Dilutions** | Flow Cyt 1:100 |
| **Purification** | Purified from cell culture supernatant by affinity chromatography |
| **Formulation & Reconstitution** | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| **Storage&Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | The protein encoded by this gene is a receptor for monocyte chemoattractant protein-1; a chemokine which specifically mediates monocyte chemotaxis. Monocyte chemoattractant protein-1 is involved in monocyte infiltration in inflammatory diseases such as rheumatoid arthritis as well as in the inflammatory response against tumors. The encoded protein mediates agonist-dependent calcium mobilization and inhibition of adenylyl cyclase. This protein can also be a coreceptor with CD4 for HIV-1 infection. This gene is located in the chemokine receptor gene cluster region of chromosome 3. |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |
| **DIMA Disclaimer** | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr |

DIMABIO CONFIDENTIAL





**Figure 1.** Flow cytometry analysis with Anti-CCR2 (DMC447) on HEK293 cells transfected with human CCR2 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com