Anti-CEACAM8 antibody(1H10), Rabbit mAb

Cat. No. DME101207

PRODUCT INFORMATION

Clone ID	**Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** 1H10
Target	CEACAM8
Synonyms	CD66b;CD67;CGM6;NCA-95
Host Species	Rabbit
Description	Anti-CEACAM8 antibody(1H10), Rabbit mAb
Delivery	In Stock
Uniprot ID	P31997
IgG type	Rabbit mAb
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1/100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Enables protein heterodimerization activity. Involved in heterophilic cell-cell adhesion via plasma membrane cell adhesion molecules. Located in cell surface and extracellular space. Biomarker of severe acute respiratory syndrome. [provided by Alliance of Genome Resources, Apr 2022]
Usage	Research use only
Conjugate	Unconjugated




Figure 1. Flow cytometry analysis with 1μg/mL Anti-CEACAM8 (1H10) mAb on HEK293 cells transfected with human CEACAM8 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com