PRODUCT INFORMATION

Clone ID	**Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** 3G1
Target	Eribulin
Synonyms	N/A
Host Species	Rabbit
Description	Anti-Eribulin antibody(3G1); Rabbit mAb
Delivery	In Stock
Uniprot ID	N/A
IgG type	Rabbit IgG
Clonality	Monoclonal
Reactivity	N/A
Applications	ELISA
Recommended Dilutions	ELISA 1/5000-10000
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	N/A
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr

DIMABIO CONFIDENTIAL




Figure 1. Elisa plates were pre-coated with IgG-Eribulin (0.2μg/per well). Serial diluted anti-Eribulin monoclonal antibody (DME101049) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Eribulin monoclonal antibody binding with IgG-Eribulin is 1.670ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com