PRODUCT INFORMATION

Common Name	**Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** RO4858696-000,teprotumumab-trbw
Conjugate	Unconjugated
Synonyms	CD221, IGFIR, IGFR, JTK13
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
Host Species	Homo sapiens
IgG type	Human IgG1(E356D,M358L) – kappa
Reactivity	Human
Target	IGF-1R
Uniprot ID	P08069
Description	Anti-IGF-1R(teprotumumab biosimilar) mAb
Delivery	In Stock
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).Lyophilized antibodies are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only

DIMABIO CONFIDENTIAL




Figure 1. IGF-1R protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with 1μg/mL Anti-IGF-1R(teprotumumab biosimilar) mAb (BME100174) on HEK293 cells transfected with human IGF-1R (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram),and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com




Figure 2. Flow cytometry analysis of antigen binding of anti-human IGF-1R mAb(BME100174).
(A) BME100174 does not bind to Jurkat cells that do not express IGF-1R.
(B) A clear peak shift of BME100174 was seen compared to the control when incubated with IGF-1R-expressing Hela cells, indicating strong binding of BME100174 to IGF-1R. Antibodies were incubated at 5 μg/mL.




Figure 3. Flow cytometry analysis of antigen binding of anti-human IGF-1R mAb(BME100174).
(A) BME100174 does not bind to Jurkat cells that do not express IGF-1R.
(B) A clear peak shift of BME100174 was seen compared to the control when incubated with IGF-1R-expressing MCF-7 cells, indicating strong binding of BME100174 to IGF-1R. Antibodies were incubated at 5 μg/mL.

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com