## PRODUCT INFORMATION

| | |
|---|---|
| **Clone ID** | **Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** 1A1 |
| **Target** | IL13 |
| **Synonyms** | IL-13;NC30 |
| **Host Species** | Rabbit |
| **Description** | Anti-IL13 antibody(1A1), IgG1 Chimeric mAb |
| **Delivery** | In Stock |
| **Uniprot ID** | P35225 |
| **IgG type** | Rabbit/Human Fc chimeric IgG1 |
| **Clonality** | Monoclonal |
| **Reactivity** | Human |
| **Applications** | ELISA |
| **Recommended Dilutions** | ELISA 1/5000-10000 |
| **Purification** | Purified from cell culture supernatant by affinity chromatography |
| **Formulation & Reconstitution** | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| **Storage&Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | This gene encodes an immunoregulatory cytokine produced primarily by activated Th2 cells. This cytokine is involved in several stages of B-cell maturation and differentiation. It up-regulates CD23 and MHC class II expression, and promotes IgE isotype switching of B cells. This cytokine down-regulates macrophage activity, thereby inhibits the production of pro-inflammatory cytokines and chemokines. This cytokine is found to be critical to the pathogenesis of allergen-induced asthma but operates through mechanisms independent of IgE and eosinophils. This gene, IL3, IL5, IL4, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL4. [provided by RefSeq, Jul 2008] |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |
| **DIMA Disclaimer** | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr |

DIMABIO-CONFIDENTIAL





Figure 1. ELISA plate pre-coated by 1 μg/ml (100 μl/well) Human IL13(35-146), hFc tagged protein can bind Rabbit anti-IL13 monoclonal antibody(clone: 1A1) in a linear range of 20-500 ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com