## PRODUCT INFORMATION

| | |
|---|---|
| **Clone ID** | **Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806**<br>11F1 |
| **Target** | IL21 |
| **Synonyms** | IL21;Za11;Interleukin-21 |
| **Host Species** | Rabbit |
| **Description** | Anti-IL21 antibody(11F1), IgG1 Chimeric mAb |
| **Delivery** | In Stock |
| **Uniprot ID** | Q9HBE4 |
| **IgG type** | Rabbit/Human Fc chimeric IgG1 |
| **Clonality** | Monoclonal |
| **Reactivity** | Human |
| **Applications** | ELISA |
| **Recommended Dilutions** | ELISA 1/5000-10000 |
| **Purification** | Purified from cell culture supernatant by affinity chromatography |
| **Formulation & Reconstitution** | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| **Storage&Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation, proliferation and activity of multiple target cells including macrophages, natural killer cells, B cells and cytotoxic T cells. Dysregulation of this gene plays a role in multiple immune-mediated diseases including lupus, psoriasis and chronic inflammatory diseases. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2011] |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |
| **DIMA Disclaimer** | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr |

DIMABIO CONFIDENTIAL





Figure 1. ELISA plate pre-coated by 1 μg/ml (100 μl/well) Human IL21 Protein can bind Rabbit anti-IL21 monoclonal antibody(clone: 11F1) in a linear range of 5-50 ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com