Biotinylated Anti-HER2 antibody(DMC369); IgG1 Chimeric mAb

Cat. No. DMC100369B

PRODUCT INFORMATION

Clone ID	**Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** DMC369
Target	Her2
Synonyms	ERBB2;CD340;HER-2:neu;HER2;MLN19;NEU;NGL;TKR1
Host Species	Rabbit
Description	Biotinylated Anti-HER2 antibody(DMC369); IgG1 Chimeric mAb
Delivery	2-3 weeks
Uniprot ID	P04626
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However; it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer; stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways; such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported; with the most common allele; Ile654-Ile655; shown here. Amplification and:or overexpression of this gene has been reported in numerous cancers; including breast and ovarian tumors. Alternative splicing results in several additional transcript variants; some encoding different isoforms and others that have not been fully characterized.
Usage	Research use only
Conjugate	Biotinylated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com