PRODUCT INFORMATION

Target	IL31
Synonyms	Interleukin-31，IL-31
Description	Recombinant Canine IL31 protein with C-terminal 6×His tag
Delivery	In Stock
Uniprot ID	C7G0W1
Expression Host	HEK293
Tag	C-6×His tag
Molecular Characterization	IL31(Ser24-Gln159) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 16.5 kDa after removal of the signal peptide. The apparent molecular mass of dIL31-His is approximately 15-25 kDa due to glycosylation.
Purity	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	IL31, which is made principally by activated Th2-type T cells, interacts with a heterodimeric receptor consisting of IL31RA (MIM 609510) and OSMR (MIM 601743) that is constitutively expressed on epithelial cells and keratinocytes. IL31 may be involved in the promotion of allergic skin disorders and in regulating other allergic diseases, such as asthma (Dillon et al., 2004 [PubMed 15184896]).[supplied by OMIM, Mar 2008]
Usage	Research use only
Conjugate	Unconjugated

DIMA BIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com




Figure 1. Canine IL31 Protein, His Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 μg/mL (100 μL/well) Canine IL31 Protein, His Tag (PME-D100004) can bind Anti-Canine IL31(lokivetmab biosimilar) mAb (BME100268) in a linear range of 0.64–80 ng/mL.

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com