Human AGTR2 full length protein-synthetic nanodisc

Cat. No. FLP100212

## PRODUCT INFORMATION

| | |
|---|---|
| **Tag** | **Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806**<br>C-Flag Tag |
| **Target** | AGTR2 |
| **Synonyms** | AT2, ATGR2, MRX88 |
| **Description** | Human AGTR2 full length protein-synthetic nanodisc |
| **Delivery** | 6~8weeks |
| **Uniprot ID** | P50052 |
| **Expression Host** | HEK293 |
| **Protein Families** | GPCR,Transmembrane,Druggable Genome, |
| **Protein Pathways** | ACE-Inhibitor pathway PharmGKB,GPCRDB Class A Rhodopsin-like,Peptide GPCRs,Apoptosis,Cancer,Endothelial Cell Biology,G-Protein Coupled Receptors Signaling Pathway, |
| **Molecular Weight** | The human full length AGTR2 protein has a MW of 41.2kDa |
| **Formulation & Reconstitution** | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for |
| **Storage&Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | The protein encoded by this gene belongs to the G-protein coupled receptor 1 family, and functions as a receptor for angiotensin II. It is an intergral membrane protein that is highly expressed in fetus and in neonates, but scantily in adult tissues, except brain, adrenal medulla, and atretic ovary. This receptor has been shown to mediate programmed cell death and this apoptotic function may play an important role in developmental biology and pathophysiology. Mutations in this gene are been associated with X-linked cognitive disability. Severe Acute Respiratory Syndrome Coronavirus (SARS-CoV) and SARS-CoV-2 infection results in down-regulation of angiotensin converting enzyme-2 (ACE2) receptors, the effects of which, triggers serious inflammatory lesions in the tissues involved, primarily in the lungs. The inflammatory reaction appears to be mediated by angiotensin II derivatives, including the angiotensin AT2 receptor which has been found to be upregulated in bronchoalveolar lavage samples from Coronavirus disease 2019 (COVID19) patients. [provided by RefSeq, Jul 2020] |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com