## PRODUCT INFORMATION

| | |
|---|---|
| **Target** | BTLA |
| **Synonyms** | BTLA;CD272 |
| **Description** | Recombinant human BTLA protein with C-terminal 6×His tag |
| **Delivery** | In Stock |
| **Uniprot ID** | Q7Z6A9 |
| **Expression Host** | HEK293 |
| **Tag** | C-6×His Tag |
| **Molecular Characterization** | Human BTLA(Lys31-Ser150) 6×His tag |
| **Molecular Weight** | The protein has a predicted molecular mass of 14.6 kDa after removal of the signal peptide.The apparent molecular mass of Human-BTLA-His is approximately 25-35 kDa due to glycosylation. |
| **Purity** | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| **Formulation & Reconstitution** | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| **Storage&Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | This gene encodes a member of the immunoglobulin superfamily. The encoded protein contains a single immunoglobulin (Ig) domain and is a receptor that relays inhibitory signals to suppress the immune response. Alternative splicing results in multiple transcript variants. Polymorphisms in this gene have been associated with an increased risk of rheumatoid arthritis. |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |

DIMA BIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com





Figure 1. Human BTLA Protein, His Tag on SDS-PAGE under reducing condition.





Figure 2. ELISA plate pre-coated by 1 μg/ml (100 μl/well) Human BTLA , His tagged protein (PME100524) can bind Anti-BTLA Neutralizing antibody BME100054 in a linear range of 16-80 ng/ml.

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com





Figure 3. Flow cytometry analysis with 15 μg/mL Human BTLA Protein, His tag (PME100524) on HEK293 cells transfected with human HVEM (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com