PRODUCT INFORMATION

Target	CEACAM5
Synonyms	CEA; CD66e
Description	Recombinant human CEACAM5(233-322) Protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P06731
Expression Host	HEK293
Tag	C-Human Fc tag
Molecular Characterization	CEACAM5(Val233-Pro322) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 35.9 kDa after removal of the signal peptide. The apparent molecular mass of CEACAM5(233-322)-hFc is approximately 35-70 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a cell surface glycoprotein that represents the founding member of the carcinoembryonic antigen (CEA) family of proteins. The encoded protein is used as a clinical biomarker for gastrointestinal cancers and may promote tumor development through its role as a cell adhesion molecule. Additionally, the encoded protein may regulate differentiation, apoptosis, and cell polarity. This gene is present in a CEA family gene cluster on chromosome 19. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2015]
Usage	Research use only
Conjugate	Unconjugated

DIMA BIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com




Figure 1. Human CEACAM5(233-322) Protein, hFc Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 μg/mL (100 μL/well) Human CEACAM5(233-322) Protein, hFc Tag (PME101572) can bind Anti-CEACAM5 antibody, IgG1 Chimeric mAb in a linear range of 0.64–80 ng/mL.

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)
Email: info@dimabio.com
Website: www.dimabio.com