## PRODUCT INFORMATION

| | |
|---|---|
| **Target** | CEACAM8 |
| **Synonyms** | CD66b;CD67;CGM6;NCA-95 |
| **Description** | Recombinant human CEACAM8 protein with C-terminal 6×His tag |
| **Delivery** | In Stock |
| **Uniprot ID** | P31997 |
| **Expression Host** | HEK293 |
| **Tag** | C-6×His Tag |
| **Molecular Characterization** | CEACAM8(Gln35-Ser319) 6×His tag |
| **Molecular Weight** | The protein has a predicted molecular mass of 32.2 kDa after removal of the signal peptide. The apparent molecular mass of CEACAM8-His is approximately 35-70 kDa due to glycosylation. |
| **Purity** | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| **Formulation & Reconstitution** | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| **Storage&Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | Cell surface glycoprotein that plays a role in cell adhesion in a calcium-independent manner (PubMed:8776764, PubMed:2022629, PubMed:11590190). Mediates heterophilic cell adhesion with other carcinoembryonic antigen-related cell adhesion molecules, such as CEACAM6 (PubMed:8776764, PubMed:2022629, PubMed:11590190). Heterophilic interaction with CEACAM8 occurs in activated neutrophils (PubMed:8776764).[UniProtKB/Swiss-Prot Function] |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |

DIMA BIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com





Figure 1. Human CEACAM8 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com