Human CNGA2-Strep full length protein-synthetic nanodisc

Cat. No. FLP120746

## PRODUCT INFORMATION

| | |
|---|---|
| **Tag** | **Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806**<br>C-Flag&Strep Tag |
| **Target** | CNGA2 |
| **Synonyms** | CNCA, CNCA1, CNG2, OCNC1, OCNCALPHA, OCNCa |
| **Description** | Human CNGA2-Strep full length protein-synthetic nanodisc |
| **Delivery** | 6~8weeks |
| **Uniprot ID** | Q16280 |
| **Expression Host** | HEK293 |
| **Protein Families** | Ion Channels: Cyclic nucleotide gated |
| **Protein Pathways** | N/A |
| **Molecular Weight** | The human full length CNGA2-Strep protein has a MW of 76 kDa |
| **Formulation & Reconstitution** | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for |
| **Storage&Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | The protein encoded by this gene represents the alpha subunit of a cyclic nucleotide-gated olfactory channel. The encoded protein contains a carboxy-terminal leucine zipper that mediates channel formation. [provided by RefSeq, Jan 2010] |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com