Human GNRHR full length protein-synthetic nanodisc

Cat. No. FLP100092

PRODUCT INFORMATION

Tag	**Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** C-Flag Tag
Target	GNRHR
Synonyms	GNRHR1; GRHR; HH7; LHRHR; LRHR
Description	Human GNRHR full length protein-synthetic nanodisc
Delivery	In Stock
Uniprot ID	P30968
Expression Host	HEK293
Protein Families	Druggable Genome, GPCR, Transmembrane
Protein Pathways	GnRH signaling pathway, Neuroactive ligand-receptor interaction
Molecular Weight	The human full length GNRHR protein has a MW of 37.7 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The receptor for type 1 gonadotropin-releasing hormone. This receptor is a member of the seven-transmembrane, G-protein coupled receptor (GPCR) family. It is expressed on the surface of pituitary gonadotrope cells as well as lymphocytes, breast, ovary, and prostate. Following binding of gonadotropin-releasing hormone, the receptor associates with G-proteins that activate a phosphatidylinositol-calcium second messenger system. Activation of the receptor ultimately causes the release of gonadotropic luteinizing hormone (LH) and follicle stimulating hormone (FSH). Defects in this gene are a cause of hypogonadotropic hypogonadism (HH). Alternative splicing results in multiple transcript variants encoding different isoforms.
Usage	Research use only
Conjugate	Unconjugated

DIMABIO CONFIDENTIAL




Figure 1. Elisa plates were pre-coated with Flag Tag GNRHR-Nanodisc (0.2μg/per well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with GNRHR-Nanodisc is 6.385ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com




Figure 2. WB analysis of Human GNRHR-Nanodisc with anti-Flag monoclonal antibody at 1/5000 dilution, followed by Goat Anti-Rabbit IgG HRP at 1/5000 dilution

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com