PRODUCT INFORMATION

Tag	**Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** C-Flag Tag
Target	MRP4
Synonyms	MOAT-B, MOATB, MRP4
Description	Human MRP4 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	O15439
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length MRP4 protein has a MW of 149.5kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for
Storage&Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MRP subfamily which is involved in multi-drug resistance. This family member plays a role in cellular detoxification as a pump for its substrate, organic anions. It may also function in prostaglandin-mediated cAMP signaling in ciliogenesis. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Sep 2014]
Usage	Research use only
Conjugate	Unconjugated

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com