Human SUCR1-Strep full length protein-synthetic nanodisc

Cat. No. FLP120471

## PRODUCT INFORMATION

| | |
|---|---|
| **Tag** | **Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806**<br>C-Flag&Strep Tag |
| **Target** | SUCR1 |
| **Synonyms** | GPR91 |
| **Description** | Human SUCR1-Strep full length protein-synthetic nanodisc |
| **Delivery** | 6~8weeks |
| **Uniprot ID** | Q9BXA5 |
| **Expression Host** | HEK293 |
| **Protein Families** | GPCR,Transmembrane,Druggable Genome, |
| **Protein Pathways** | GPCRDB Class A Rhodopsin-like, |
| **Molecular Weight** | The human full length SUCR1-Strep protein has a MW of 38.7 kDa |
| **Formulation & Reconstitution** | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for |
| **Storage&Shipping** | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| **Background** | This gene encodes a G-protein-coupled receptor for succinate, an intermediate molecule of the citric acid cycle. It is involved in the promotion of hematopoietic progenitor cell development, and it has a potential role in renovascular hypertension which has known correlations to renal failure, diabetes and atherosclerosis. [provided by RefSeq, Oct 2009] |
| **Usage** | Research use only |
| **Conjugate** | Unconjugated |

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA)
/+86-18062749453(China)
/+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com