PE-conjugated Anti-MICB antibody(DM168); Rabbit mAb

Cat. No. DME100168P

## PRODUCT INFORMATION

| | |
|---|---|
| **Clone ID** | **Warning**: Undefined variable $hasAttributeValueDescription in **/www/wwwroot/tech.dimabio.cn/wp-content/plugins/woocommerce-print-products/public/class-woocommerce-print-products-public.php** on line **2806** DM168 |
| **Target** | MICB |
| **Synonyms** | MIC-B; PERB11.2 |
| **Host Species** | Rabbit |
| **Description** | PE-conjugated Anti-MICB antibody(DM168); Rabbit mAb |
| **Delivery** | Under Development |
| **Uniprot ID** | Q29980 |
| **IgG type** | Rabbit IgG |
| **Clonality** | Monoclonal |
| **Reactivity** | Human |
| **Applications** | Flow Cyt |
| **Recommended Dilutions** | Flow Cyt 1:100 |
| **Purification** | Purified from cell culture supernatant by affinity chromatography |
| **Formulation & Reconstitution** | Liquid□PBS with 0.05% Proclin300, 1% BSA |
| **Storage&Shipping** | Store at 2°C-8°C for 6 months |
| **Background** | This gene encodes a heavily glycosylated protein which is a ligand for the NKG2D type II receptor. Binding of the ligand activates the cytolytic response of natural killer (NK) cells; CD8 alphabeta T cells; and gammadelta T cells which express the receptor. This protein is stress-induced and is similar to MHC class I molecules; however; it does not associate with beta-2-microglobulin or bind peptides. Alternative splicing results in multiple transcript variants. |
| **Usage** | Research use only |
| **Conjugate** | PE-conjugated |
| **DIMA Disclaimer** | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scr |

DIMABIO CONFIDENTIAL

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com
Website: www.dimabio.com